



# Sal I

Cat # RE115

Pack Size: 500U/100uL(5U/uL)

Storage: -20°C

Recognition Sequence:  $5' \text{GTCGAC} 3'$   
 $3' \text{CAGCTG} 5'$

Optimal Buffer: 10x Universal Buffer □ 1mL

## Introduction

Sal I is a restriction endonuclease that recognizes and cleaves the specific 5'-GTCGAC-3' DNA sequence. This enzyme is widely used in molecular biology for cloning, mapping, and DNA manipulation.

## Features

- Assayed on λ DNA
- Heat inactivation: 65°C for 20 minutes
- Ligation/recutting assay: After 20-fold overdigestion with EcoRV, >90% of the DNA fragments can be ligated and recut
- Overdigestion assay: No nonspecific activity detected after incubation of 1 μg of λ DNA with 20 units of EcoRV for 16 hours

## Protocol

- Reaction setup (This is just an example to show the relative concentrations and volumes in the reaction may wish to set up a reaction ranging from 10 μl to 200 μl or more

Component	50 μl Reaction
DNA	1 μg
10x Universal Buffer	5 μl
Sal I	2-5 units
Nuclease-free Water	to 50μl

- Incubate at 37°C for 10 minutes. Longer incubation times (sometimes overnight) may be followed as per digestion efficiency
- Heat inactivate enzyme at 65°C for 10 mins.
- Please note that supercoiled plasmid DNA and PCR fragments may have varied rate of cleavage and sometime needs more time to completely digest

## Certificate of Analysis

Source	:An <i>E.coli</i> strain, that carries the cloned Sal I gene from <i>Streptomyces albus</i> .
Supplied in	:10mM Tris-Hcl (pH 7.6), 50mM NaCl, 0.1mM EDTA, 1mM DTT, 100ug/ml, BSA, 50% Glycerol.
Reaction Conditions	:1x Universal Buffer, Incubate at 37°C for 10 min
Unit definition	:One unit of Sal I is defined as the amount of enzyme required to completely digest 1 μg of lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.
Optimal temperature	:37°C
Heat Inactivation	:Enzyme is inactivated by incubation at 65°C for 10 minutes.

## Quality Control Assays

### Ligation of DNA fragments

DNA fragments are produced by an excessive over digestion of substrate DNA with each restriction endonuclease. These fragments are then ligated with T4 DNA Ligase at a 5' termini concentration of 0.1-1.0 μM. The ligated fragments are then recut with the same restriction endonuclease. Ligation can only occur if the 3' and 5' termini are left intact, and only those molecules with a perfectly restored recognition site can be recleaved. A normal banding pattern after cleavage indicates that both the 3' and 5' termini are intact, and the enzyme preparation is free of detectable exonucleases and phosphatases.

### DNA digestion with Sal I may be affected by some types of methylation

\* In general, it is recommended to use 5–10 units of enzyme per μg of plasmid DNA, and 10–20 units for genomic DNA in a 1-hour digest. Enzyme volume should not exceed 10% of the total reaction volume to prevent star activity.

Reagents Supplied with Enzyme: 10x Universal Buffer